

## SCIENTIFIC NOTE

### FIRST RECORD OF INVASIVE MOSQUITO *Aedes albopictus* IN TABASCO AND YUCATAN, MEXICO

ALDO I. ORTEGA-MORALES,<sup>1</sup> GUILLERMO BOND,<sup>2</sup> RAMÓN MÉNDEZ-LÓPEZ,<sup>1</sup>  
JAVIER A. GARZA-HERNÁNDEZ,<sup>3</sup> LUIS M. HERNÁNDEZ-TRIANA<sup>4</sup> AND MAURICIO CASAS-MARTÍNEZ<sup>2,5</sup>

**ABSTRACT.** The invasive mosquito *Aedes albopictus* is currently distributed in most of the southern Mexican region. Since the species was first recorded in the state of Tamaulipas, in northeastern Mexico in 1988, it has expanded its distribution throughout the Sierra Madre Oriental and Gulf of Mexico to the Neotropical region of the country. Currently the species occurs in the states of Tamaulipas, Coahuila, Nuevo Leon, Veracruz, Chiapas, Morelos, Quintana Roo, Sinaloa, San Luis Potosi, and Hidalgo. This is the first report of the mosquito in the states of Tabasco and Yucatan and the confirmation of its presence in Quintana Roo state. *Aedes albopictus* has been incriminated as a secondary vector of diseases such as those caused by dengue, chikungunya, and Zika viruses, which have caused epidemic outbreaks in most tropical and subtropical regions of Mexico; therefore, surveillance for the detection of *Ae. albopictus* is paramount so that targeted control strategies can be implemented for its control throughout Mexico.

**KEY WORDS** *Aedes albopictus*, first record, Mexico, Quintana Roo, Tabasco, Yucatan

The Asian tiger mosquito, *Aedes (Stegomyia) albopictus* (Skuse), is the most invasive mosquito species worldwide (GISD 2017). *Aedes albopictus* has a Pantropical distribution and has been reported in 81 countries (WRBU 2005). The introduction of *Ae. albopictus* into the American continent occurred in the 20th century, where it was detected for the first time in the USA in Harris County, Texas, breeding in discarded tires imported from northern Asia (Sprenger and Wuithiranyagool 1986, Hawley et al. 1987, Reiter and Sprenger 1987, Craven et al. 1988), and it rapidly adapted to the Nearctic environmental conditions of the southern United States. This species was detected for first time in Mexico in the northeastern state of Tamaulipas in 1988 (Francy et al. 1990) and since then has had a rapid and successful dispersion in almost all tropical and subtropical regions of the country. Currently *Ae. albopictus* has been reported in the states of Tamaulipas (Francy et al. 1990, Ibáñez-Bernal et al. 1997, Reyes-Villanueva et al. 2013, Ortega-Morales et al. 2015), Coahuila (Ibáñez-Bernal and

Martínez-Campos 1994, Rodríguez and Ortega 1994), Nuevo Leon (Pesina et al. 2001, Orta-Pesina et al. 2005, Reyes-Villanueva et al. 2013), Veracruz (Flisser et al. 2002), Chiapas (Casas-Martínez and Torres-Estrada 2003), Quintana Roo (Salomón-Gracia et al. 2012), Morelos (Villegas-Trejo et al. 2010), Sinaloa (Torres et al. 2015), San Luis Potosí (Ortega-Morales and Siller-Rodríguez 2016), and Hidalgo (Ortega-Morales et al. 2016) (Fig. 1).

In Mexico, *Aedes aegypti* (L.) and *Ae. albopictus* are the most important mosquitoes responsible for the transmission of arboviruses such as dengue (DENV), chikungunya (CHIKV), and Zika (ZIKV). Consequently the distribution and colonization of those species in new regions are systematically monitored by personnel from the Mexican vector-borne disease surveillance and control programs (SEGOB 2015) and many other research groups. To update the checklist of mosquito species in the southeastern states of Mexico, which include the Yucatan Peninsula region (states of Campeche, Yucatan, and Quintana Roo) and Tabasco state, mosquitoes in their different stages of life (larvae, pupae, and adults) were collected during the dry and rainy seasons of 2015–17. The collecting methodology consisted of the search for immature stages of mosquitoes in all available breeding sites, such as artificial containers (vases, bottles, buckets, discarded tires) and natural containers (bamboo internodes, tree holes, axils of plants). In addition, ovitraps (a black plastic container of 1-liter capacity filled with potable water with germination paper strips on the inner edge of the water, where gravid females lay eggs) were also used in the study areas together with Centers for Disease Control and Prevention (CDC) light traps, a BDV tent trap, and landing/biting catches. All collected

<sup>1</sup> Departamento de Parasitología, Universidad Autónoma Agraria Antonio Narro, Unidad Laguna, Torreón, Coahuila, Mexico 27084.

<sup>2</sup> Centro Regional de Investigación en Salud Pública, Instituto Nacional de Salud Pública, Tapachula, Chiapas, Mexico 30700.

<sup>3</sup> Laboratorio de Biotecnología, Instituto de Ciencias Biomédicas, Universidad Autónoma de Ciudad Juárez, Chihuahua, Mexico 32310.

<sup>4</sup> Animal and Plant Health Agency, Department of Virology, Wildlife Zoonoses and Vector Borne Diseases Research Group, Woodham Lane, New Haw, Addlestone, Surrey KT153NB, United Kingdom.

<sup>5</sup> To whom correspondence should be addressed.




Fig. 1. Distribution of *Aedes albopictus* in Tabasco state and the Yucatan Peninsula.

specimens were transported alive to the Molecular Biology Laboratory of the Agrarian Autonomous University Antonio Narro Laguna Unit (UAAAN-UL). Adult mosquitoes were killed in lethal chambers and kept in nitrogen vapor until identification. Immature stages were reared individually to obtain larval and pupal skins. The ovitraps strips were transported to the Regional Center for Public Health Research (CIRSP-INSP) and placed in trays filled with water for emergence. All specimens were identified using available taxonomic keys. All material was deposited in the Culicidae Collection of UAAAN-UL (accession numbers CUA1414-B and CUAC1414-C) and in the Biological Collection of Mosquitoes of Medical Importance at CIRSP-INSP.

The present study reports the presence of *Ae. albopictus* in 4 municipalities in Tabasco (Huimanguillo, Teapa, Tacotalpa, and Villahermosa), 3 in Quintana Roo (Cancun, Felipe Carrillo Puerto, and Tulum), and 1 in Yucatan (Tizimin) (Table 1). This is the first record of *Ae. albopictus* in the states of Tabasco and Yucatan; so far, it has not been detected in the state of Campeche. Since the discovery of the presence of *Ae. albopictus* in Mexico, the species has widened its distribution in Nearctic and Neotropical

regions of the southeastern part of the country, and it is now established in many physiographic regions, such as the Sierra Madre Oriental, the coastal plain of the North and South Gulf, the Yucatan Peninsula, and the Sierra Madre of the South. In these regions, *Ae. albopictus* finds favorable environmental conditions for its development, such as abundant vegetation, a humid and warm climate, and great availability of natural and artificial breeding sites within human settlements. Some cities on the Yucatan Peninsula, such as Cancun, Tulum, Playa del Carmen, and Chichen-Itza, are some of the main tourist destinations in Mexico. Therefore, the probable presence of *Ae. albopictus* in these cities could represent an important risk of infection of DENV, CHIKV, and ZIKV for domestic and international travelers.

Because of its medical importance, we recommend widening the surveillance of this species to other regions of Mexico, especially the Pacific coast (states of Guerrero and Oaxaca) and along the southern border.

We thank Salvador Morales-Avitia and Martha Ortega-Lozano for their valuable collaboration during field collections trips in Tabasco state and Miguel Muñoz-Reyes, José L. Aguilar-Rodriguez, and Mar-

Table 1. Records of *Aedes albopictus* in Tabasco, Yucatan, and Quintana Roo states, Mexico

Collection Date	Location	State	Longitude N	Latitude W	Elevation	Habitat
July 18, 2015	Huimanguillo	Tabasco	17°34'11.6"	−93°21'27"	72 m	Flower vase
July 18, 2015	Huimanguillo	Tabasco	17°34'14"	−93°31'23"	72 m	Discarded tire
July 18, 2015	Huimanguillo	Tabasco	17°36'16"	−93°31'13"	68 m	Discarded tire
October 19, 2015	Huimanguillo	Tabasco	17°55'19"	−93°46'12"	17 m	Plastic bottle
October 20, 2015	Huimanguillo	Tabasco	17°50'14"	−93°55'12"	22 m	Plastic bottle
October 20, 2015	Huimanguillo	Tabasco	17°24'36"	−93°37'53"	59 m	Plastic drum
October 20, 2015	Huimanguillo	Tabasco	17°50'14"	−93°55'12"	22 m	Intradomiciliary
October 20, 2015	Huimanguillo	Tabasco	17°48'50"	−93°52'18"	10 m	Bamboo internode
October 20, 2015	Huimanguillo	Tabasco	17°45'5"	−93°42'27"	55 m	Discarded tire
October 20, 2015	Huimanguillo	Tabasco	17°45'55"	−93°4'9"	—	Tree hole
October 20, 2015	Huimanguillo	Tabasco	17°40'47"	−93°30'15"	68 m	Biting
October 20, 2015	Huimanguillo	Tabasco	17°46'38"	−93°38'21"	41 m	Biting
October 20, 2015	Huimanguillo	Tabasco	17°40'48"	−93°30'16"	—	Tree hole
October 20, 2015	Huimanguillo	Tabasco	17°40'1"	−93°28'53"	—	Biting
October 20, 2015	Huimanguillo	Tabasco	17°39'42"	−93°28'12"	54 m	Tree hole
October 21, 2015	Huimanguillo	Tabasco	17°20'47"	−93°35'33"	174 m	Discarded tire
October 21, 2015	Huimanguillo	Tabasco	17°23'29"	−93°36'39"	184 m	<i>Xanthosoma</i> axil
October 21, 2015	Huimanguillo	Tabasco	17°24'36"	−93°37'53"	59 m	CDC light trap
October 22, 2015	Huimanguillo	Tabasco	17°22'44"	−93°38'23"	515 m	Biting
October 27, 2015	Teapa	Tabasco	17°31'10"	−92°55'31"	79 m	Flower vase
October 27, 2015	Teapa	Tabasco	17°30'14"	−92°53'57"	117 m	Plastic bottle
October 29, 2015	Tacotalpa	Tabasco	17°30'14"	−92°46'53"	39 m	Metal can
October 30, 2015	Tacotalpa	Tabasco	17°26'17"	−92°44'1"	405 m	Coconut shell
October 8, 2016	Villahermosa	Tabasco	17°57'21"	−92°59'15"	8 m	Glass
October 9, 2016	Villahermosa	Tabasco	17°57'36"	−92°57'25"	14 m	Cement vase
November 7, 2016	Villahermosa	Tabasco	17°55'30"	−93°01'42"	20 m	Aluminum can
November 7, 2016	Villahermosa	Tabasco	17°54'19"	−93°02'12"	18 m	Discarded tire
February 1, 2017	Villahermosa	Tabasco	17°57'21"	−92°59'15"	8 m	Human biting
October 21, 2016	Cancun	Quintana Roo	20°08'33"	−86°52'49"	7 m	Variety of vases
February 10, 2017	Felipe Carrillo Puerto	Quintana Roo	20°04'35"	−87°37'05"	6 m	Rock hole
February 11, 2017	Felipe Carrillo Puerto	Quintana Roo	20°04'35"	−87°37'05"	6 m	Variety of vases
February 12, 2017	Tulum	Quintana Roo	20°12'25"	−87°28'08"	19 m	Disposable cup
February 13, 2017	Cancun	Quintana Roo	20°08'33"	−86°52'49"	7 m	Plastic bottle
June 19, 2017	Tizimin	Yucatan	21°30'51"	−87°41'58"	3 m	BDV tent trap
June 20, 2017	Tizimin	Yucatan	21°30'51"	−87°41'58"	3 m	Ovitrap

tin Vázquez-Castillo for their support on collection trips in the Yucatan Peninsula. We also thank the Secretary of Public Education (SEP-PRODEP) and the National Commission for the Knowledge and Use of Biodiversity (CONABIO) for providing funds to carry out the field trips of this study project: “Taxonomic inventory of some groups of haematophagous nematoceroids (Lower Diptera) in the state of Tabasco, Mexico” (Grant No. CUAC1414) and “Diagnosis of the current status of biological invasion of *Aedes albopictus* in the main protected natural areas of the Yucatan Peninsula” (Grant No. LI040).

REFERENCES CITED

Casas-Martinez M, Torres-Estrada JL. 2003. First evidence of *Aedes albopictus* (Skuse) in Southern Chiapas, Mexico. *Emer Infect Dis* 9:606–607.

Craven RB, Eliason DA, Francy DB, Reiter P, Campos EG, Lakob WL. 1988. Importation of *Aedes albopictus* and other exotic mosquito species into the United States in used tires from Asia. *J Am Mosq Control Assoc* 4:138–142.

Flisser A, Velasco-Villa A, Martínez-Campos C, González-Domínguez F, Briseño-García B, García R, Caballero-Servín A, Hernández-Monroy I, García-Lozano H, Gutiérrez-Cogco L, Rodríguez-Ángeles G, López-Martínez I, Galindo-Virgen S, Vázquez-Campuzano R, Balandrano-Campos S, Guzmán-Bracho C, Olivo-Díaz A, de la Rosa JL, Magos C, Escobar-Gutiérrez A, Correa D. 2002. Infectious diseases in Mexico: a survey from 1955–2000. *Arch Med Res* 33:343–350.

Francy DB, Moore CG, Eliason DA. 1990. Past, present and future of *Aedes albopictus* in the United States. *J Am Mosq Control Assoc* 1:127–132.

GISD [Global Invasive Species Database]. 2017. *Species profile: Aedes albopictus* [Internet]. Available from: <http://www.iucngisd.org/gisd/speciesname/Aedes+albopictus> [accessed October 15, 2017].

Hawley WA, Reiter P, Copeland RW, Pumpini CB, Craig GB. 1987. *Aedes albopictus* in North America: probable introduction in used tires from northern Asia. *Science* 236:1114–1116.

Ibáñez-Bernal S, Briseño B, Mutebi JP, Argot A, Rodríguez G, Martínez-Campos C, Perez R, de la Fuente-San Roman P, Tapia-Conyer R, Flisser A. 1997. First record in America of *Aedes albopictus* naturally infected with dengue virus during the 1995 outbreak at Reynosa, Mexico. *Med Vet Entomol* 11:305–309.

- Ibáñez-Bernal S, Martínez-Campos C. 1994. *Aedes albopictus* in Mexico. *J Am Mosq Control Assoc* 10:231–232.
- Orta-Pesina H, Mercado-Hernández R, Elizondo-Leal JF. 2005. Distribución de *Aedes albopictus* (Skuse) en Nuevo León, México, 2001–2004. *Salud Pub Mex* 47:163–165.
- Ortega-Morales AI, Cueto-Medina SM, Rodríguez QK. 2016. First record of the Asian tiger mosquito *Aedes albopictus* in Hidalgo state, Mexico. *J Am Mosq Control Assoc* 32:234–236.
- Ortega-Morales AI, Siller-Rodríguez QK. 2016. First record of *Aedes albopictus* (Diptera: Culicidae) in San Luis Potosí, Mexico. *J Vector Ecol* 41:314–315.
- Ortega-Morales A, Zavortink TJ, Huerta-Jiménez H, Sánchez-Ramos FJ, Valdés-Perezgasca MT, Reyes-Villanueva F, Siller-Rodríguez QK, Fernández-Salas I. 2015. Mosquito records from Mexico: the mosquitoes (Diptera: Culicidae) of Tamaulipas state. *J Med Entomol* 52:171–184.
- Pesina OH, Mercado R, Valdez-Rodríguez MA. 2001. *Aedes albopictus* in Allende City, Nuevo León, Mexico. *J Am Mosq Control Assoc* 17:260–261.
- Reiter P, Sprenger D. 1987. The used tire trade: a mechanism for the worldwide dispersal of container breeding mosquitoes. 1987. *J Am Mosq Control Assoc* 44:396–399.
- Reyes-Villanueva F, Garza-Hernández JA, García-Munguía AM, Howard AF, Ortega-Morales AI, Adeleke MA, Rodríguez-Pérez MA. 2013. *Aedes albopictus* in north-east Mexico: an update of adult distribution and first report of parasitism by *Ascogregarina taiwanensis*. *J Vector Borne Dis* 50:202–205.
- Rodríguez ML, Ortega MG. 1994. *Aedes albopictus* in Muzquiz city, Coahuila, Mexico. *J Am Mosq Control Assoc* 10:587.
- Salomón-Grajales J, Lugo-Moguel GV, Tinal-Gordillo VR, de la Cruz-Velázquez J, Beaty BJ, Eisen L, Lozano-Fuentes S, Moore CG, García-Rejón JE. 2012. *Aedes albopictus* mosquitoes, Yucatan Peninsula, Mexico. *Emerg Infect Dis* 18:525–527.
- SEGOB [Secretaría de Gobernación]. 2015. *Norma Oficial Mexicana NOM-032-SSA2-2014, Para la vigilancia epidemiológica, promoción, prevención y control de las enfermedades transmitidas por vectores*, as amended April 16, 2015. Diario Oficial de la Federación, Mexico.
- Sprenger D, Wuithiranyagool T. 1986. The discovery of *Aedes albopictus* in Harris County, Texas. *J Am Mosq Control Assoc* 2:217–219.
- Torres-Avendaño JI, Castillo-Ureta H, Torres-Montoya EH, Meza-Carrillo E, Lopez-Mendoza RL, Vazquez-Martinez MG, Rendon-Maldonado JG. 2015. First record of *Aedes albopictus* in Sinaloa, Mexico. *J Am Mosq Control Assoc* 2:164–166.
- Villegas-Trejo A, Manrique-Saide P, Che-Mendoza A, Cruz-Canto W, Fernández MG, González-Acosta C, Dzul-Manzanilla F, Huerta H, Arredondo-Jiménez JI. 2010. First report of *Aedes albopictus* and other mosquito species in Morelos, Mexico. *J Am Mosq Control Assoc* 3:321–323.
- WRBU [Walter Reed Biosystematics Unit]. 2005. *Systematic catalog of Culicidae* [Internet]. Suitland, MD: WRBU [accessed October 15, 2017]. Available from: [www.mosquitocatalog.org](http://www.mosquitocatalog.org).